

١/٥٨

ملحق رقم ٢ -

٢. معايير بذور الخضار

I - Seed standards for Vegetable seeds**I - 1- Analytical parameters**

Analytical parameters as defined by the International Seed Testing Association (ISTA rules)			Scientific name	SPECIES
Minimum analytical purity (% by weight)	Maximum content of seeds of other plant species (% by weight)	Minimum germination (% clusters or pure seed)		
96	0,5	70	<i>Asparagus officinalis</i>	Asparragus
98	0,1	75	<i>Phaseolus vulgaris</i>	Bean
95	1	70	<i>Scorzonera hispanica</i>	Black Salsify
97	1	75	<i>Brassica oleracea</i>	Broccoli,
97	1	75	<i>Brassica oleracea</i>	Brussels sprouts
97	1	75	<i>Brassica oleracea</i>	Cabbage
96	0,5	65	<i>Cynara cardunculus</i>	Cardoon
95	1	65	<i>Daucus carota</i>	Carrot
97	1	70	<i>Brassica oleracea</i>	Cauliflower
97	1	70	<i>Apium graveolens</i>	Celery
97	1	75	<i>Brassica pekinensis</i>	Chinese cabbage
98	0,1	80	<i>Cucumis sativus</i>	Cucumber
96	0,5	65	<i>Solanum melongena</i>	Eggplant
95	1	65	<i>Cichorium endivia</i>	Endive
98	0,1	80	<i>Vicia faba</i>	Faba bean
96	1	70	<i>Foeniculum vulgare</i>	Fennel
97	1	75	<i>Brassica oleracea</i>	Kohlrabi
97	0,5	65	<i>Allium porrum</i>	Leek
95	0,5	75	<i>Lactuca sativa</i>	Lettuce
98	0,1	75	<i>Cucurbita pepo</i>	Marrow,
98	0,1	75	<i>Cucumis melo</i>	Melon
97	0,5	70	<i>Allium cepa</i>	Onion
97	1	65	<i>Petroselinum crispum</i>	Parsley
98	0,1	80	<i>Pisum sativum</i>	Pea
97	0,5	65	<i>Capsicum annum</i>	Pepper
98	0	90	<i>Zea mays</i>	Pop corn
97	1	70	<i>Raphanus sativus</i>	Radish
97	0,5	70	<i>Beta vulgaris</i>	Red Beet
97	1	75	<i>Brassica oleracea</i>	Red cabbage
97	1	75	<i>Brassica oleracea</i>	Savoy cabbage
97	1	75	<i>Spinacia oleracea</i>	Spinach
98	0	90	<i>Zea mais</i>	Sweet corn
97	0,5	75	<i>Lycopersicon lycopersicum</i>	Tomato
97	1	80	<i>Brassica rapa</i>	Turnip
98	0,1	75	<i>Citrullus lanatus</i>	Water melon

III- Seed standards for fodder beet and sugar beet seeds

III - 1 - Analytical parameters

Germination parameters			Analytical purity			SEED	SPECIES
Maximum percentage of clusters giving 3 or more seedlings	Minimum percentage of germinated clusters giving a single seedling	Minimum germination (% clusters or pure seed)	Minimum analytical purity (% by weight)	Maximum content of seeds of other plant species (% by weight)	Maximum moisture content (% by weight)		
5	90	80	97	0.3	15	Monogerm seed	SUGAR BEET
5	70	75	97	0.3	15	Precision seed	
-	-	73	97	0.3	15	Natural seed of varieties with more than 85% diploids	
-	-	68	97	0.3	15	Natural seed of varieties with 15 % or more triploids and tetraploids	
-	-	73	97	0.3	15	Natural seed of varieties with more than 85% diploids	
5	58	73	97	0.3	15	Precision seed of varieties with more than 85% diploids	
-	-	68	97	0.3	15	Natural seed of varieties with 15 % or more triploids and tetraploids	FODDER BEET
5	63	68	97	0.3	15	Precision seed of varieties with 15 % or more triploids and tetraploids	

II - Seed standards for forage and grasses seeds

II - 1 Analytical parameters

Germination as defined by ISTA rules		Analytical purity as defined by ISTA rules							SPECIES				
Minimum germination (% pure seed)	Maximum content of hard seed (% of pure seed)	Minimum analytical purity (% by weight)	Maximum content of seeds of other plant species (% by weight)			Maximum content of seeds of other plant species in a sample of the weight specified in column A below							
			Total	A single species	<i>Agropyron repens</i>	<i>Alopecurus myosuroides</i>	<i>Meilolus spp</i>	Column A Weight in g		Avena fatua, Avena ludoviciana, Avena sterilis	Cuscuta spp	Rumex spp other than R. acetosella and R. maritimus	
80	40	97	1.5	1				0.3	50	0	0	10	Medicago sativa
80	5	98	0.5	0.3				0.3		0	0	5	Vicia faba
75		96	1.5	1	0.5	0.3			60	0	0	5	Lolium multiflorum
80		96	1.5	1	0.5	0.3			60	0	0	5	Lolium perenne
80		95	1.5	1	0.5	0.3			50	0	0	5	Festuca arundinacea
75		85	2	1	0.3	0.3			1000	0	0	2	Poa pratensis